

Vol. 18, No. 4, December, 2019

ISSN (Print): 0972-6268; ISSN (Online) : 2395-3454

NATURE ENVIRONMENT & POLLUTION TECHNOLOGY

*A Multidisciplinary, International Journal
on Diverse Aspects of Environment*



Technoscience Publications

website: www.neptjournal.com



Technoscience Publications

A-504, Bliss Avenue, Balewadi,
Opp. SKP Campus, Pune-411 045
Maharashtra, India

www.neptjournal.com

Nature Environment and Pollution Technology

(An International Quarterly Scientific Research Journal)

EDITORS

Dr. P. K. Goel

Former Head, Deptt. of Pollution Studies
Y. C. College of Science, Vidyanagar
Karad-415 124, Maharashtra, India

Dr. K. P. Sharma

Former Professor, Deptt. of Botany
University of Rajasthan
Jaipur-302 004, India

Published by : Mrs. T. P. Goel, B-34, Dev Nagar, Tonk Road, Jaipur-302 018
Rajasthan, India

Managing Office : Technoscience Publications, A-504, Bliss Avenue, Balewadi,
Pune-411 045, Maharashtra, India

E-mail : contact@neptjournal.com; operationst@neptjournal.com
journalnept@gmail.com

INSTRUCTIONS TO AUTHORS

Scope of the Journal

The Journal publishes original research/review papers covering almost all aspects of environment like monitoring, control and management of air, water, soil and noise pollution; solid waste management; industrial hygiene and occupational health hazards; biomedical aspects of pollution; conservation and management of resources; environmental laws and legal aspects of pollution; toxicology; radiation and recycling etc. Reports of important events, environmental news, environmental highlights and book reviews are also published in the journal.

Format of Manuscript

- The manuscript (*mss*) should be typed in double space leaving wide margins on both the sides.
- First page of *mss* should contain only the title of the paper, name(s) of author(s) and name and address of Organization(s) where the work has been carried out along with the affiliation of the authors.

Continued on back inner cover...

... Continued from inner front cover

- The text of the manuscript should run into **Abstract, Introduction, Materials & Methods, Results, Discussion, Acknowledgement** (if any) and **References** or other suitable headings in case of reviews and theoretically oriented papers. However, short communication can be submitted in running with **Abstract and References**. The references should be in full with the title of the paper.
- The figures should preferably be made on a computer with high resolution and should be capable of withstanding a reasonable reduction with the legends provided separately outside the figures. Photographs may be black and white or colour.
- Tables should be typed separately bearing a short title, preferably in vertical form. They should be of a size, which could easily be accommodated in the page of the Journal.
- References in the text should be cited by the authors' surname and year. In case of more than one reference of the same author in the same year, add suffix a,b,c,... to the year. For example: (Thomas 1969, Mass 1973a, 1973b, Madony et al. 1990, Abasi & Soni 1991).

List of References

The references cited in the text should be arranged alphabetically by authors' surname in the following manner: (Note: The titles of the papers should be in running 'sentence case', while the titles of the books, reports, theses, journals, etc. should be in 'title case' with all words starting with CAPITAL letter.)

- Dutta, A. and Chaudhury, M. 1991. Removal of arsenic from groundwater by lime softening with powdered coal additive. J. Water Supply Res. Techno. Aqua., 40(1) : 25-29.
- Hammer, D.A. (ed.) 1989. Constructed Wetlands for Wastewater Treatment-Municipal, Industrial and Agricultural. Lewis Publishers Inc., pp. 831.
- Haynes, R. J. 1986. Surface mining and wetland reclamation. In: Harper, J. and Plass, B. (eds.) New Horizons for Mined Land Reclamation. Proceedings of a National Meeting of the American Society for Surface Reclamation, Princeton, W.V.

Submission of Papers

- The paper can be submitted by e-mail as an attachment in a single WORD file at **contact@neptjournal.com**
- The paper can also be submitted online in a single WORD file through the journal's website: **www.neptjournal.com**

Attention

1. Any change in the authors' affiliation may please be notified at the earliest.
2. Please make all the correspondence by e-mail, and authors should always quote the manuscript ID (number).

Note: In order to speed up the publication, authors are requested to send the publication charges as soon as they get the 'initial acceptance' letter, and also correct the galley proof immediately after receipt. The galley proof must be checked with utmost care, as publishers owe no responsibility for mistakes. The papers will be put on priority for publication only after receiving the processing and publication charges.

Nature Environment and Pollution Technology

(Abbreviation: Nat. Env. Poll. Tech.)

(An International Quarterly Scientific Journal)

Published by



Technoscience Publications

A-504, Bliss Avenue, Opp. SKP Campus
Balewadi, Pune-411 045, Maharashtra, India

In association with

Technoscience Knowledge Communications

Mira Road, Mumbai, India

For further details of the Journal please visit the website. All the papers published on a particular subject/topic or by any particular author in the journal can be searched and accessed by typing a keyword or name of the author in the 'Search' option on the Home page of the website. All the papers containing that keyword or author will be shown on the home page from where they can be directly downloaded.

www.neptjournal.com

©Technoscience Publications: The consent is hereby given that the copies of the articles published in this Journal can be made only for purely personal or internal use. The consent does not include copying for general distribution or sale of reprints.

Published for Proprietor, Printer and Publisher: Mrs. T. P. Goel, B-34, Dev Nagar, Tonk Road, Jaipur, Rajasthan, India; Editors: Dr. P. K. Goel and Prof. K. P. Sharma

Nature Environment and Pollution Technology

Vol. 18, No. (4), December, 2019

CONTENTS

1. **Sabhan, Alan Frendy Koropitan, Mulia Purba and Widodo Setiyo Pranowo**, 3D Simulation Model of Tidal, Internal Mixing and Turbulent Kinetic Energy of Palu Bay 1083-1094
2. **Ali Abid Abojassim and Leith Hani Rasheed**, Mapping of Terrestrial Gamma Radiation in Soil Samples at Baghdad Governorate (Karakh Side), Using GIS Technology 1095-1106
3. **Yuan Ren, Dongmei Wang and Xiaochen Li**, Impacts of Human Disturbances on Riparian Herbaceous Communities in a Chinese Karst River 1107-1118
4. **Vani Sharma and Padma Singh**, Biosorption of Chromium by *Bacillus subtilis* Isolated from Ganga River 1119-1129
5. **Yuzheng Lv, Jihao Zhou, Zhengjun Mai and Jie Liu**, Nano-porous Membrane Process for Brackish Groundwater Treatment: Efficiency Analysis Using Response Surface Methodology 1131-1141
6. **Xin Yu, Haoming Fan, Liquan Liu and Jing Chen**, Calculation of Water Volume for Sediment Transport in the Sediment-laden River of the Main Stream of Liaohe River, Northern China 1143-1156
7. **Mingming Du, Fengchao Wang, Jinling Li, Tao Yu and Chengtun Qu**, Control of Air Pollutants Emission and Improvement of Incineration Rate During Incineration of Oily Sludge-Based Briquette 1157-1167
8. **Muhammad A. Asadi, Yody A.P. Ritonga, Defri Yona and Asus M.S. Hertika**, Vertical Distribution of Microplastics in Coastal Sediments of Bama Resort, Baluran National Park, Indonesia 1169-1176
9. **Deepak Singh and Gunjan Goswami**, Diversity and Seasonal Variations of Plankton Communities After Major Flash Flood in the River Mandakini of Garhwal Himalaya 1177-1184
10. **Keerthan Krishna, Krishna Murthy and G.T. Mahesha**, Performance Studies on Non-Edible Oil as IC Engine Fuels: A Review 1185-1192
11. **Fatima Zahra Hafiane, Latifa Tahri, Najia Ameer, Rajaa Rochdi, Karim Arifi and Mohammed Fekhaui**, Antibiotic Resistance of *Pseudomonas aeruginosa* in Well Waters in Irrigated Zone (Middle Atlas-Morocco) 1193-1200
12. **Yuanyuan Cai, Yaowei Du, Yue Wang, Jun Song, Bing Liu, Chenhu Zhang and Muqing Qiu**, Adsorption of Copper Ions in Aqueous Solution by Montmorillonite-Biochar Composite 1201-1209
13. **Qingkong Cai, Erjun Li, Jiechen Pan and Chao Chen**, Retrieval of the Canopy Chlorophyll Density of Winter Wheat from Canopy Spectra Using Continuous Wavelet Analysis 1211-1218
14. **Venkatesan Rajasekar, Varuvel Edwin Geo, Leenus Jesu Martin and Beddhanan Nagalingam**, Simultaneous Reduction of Smoke and NO Emission Using Lower Order Alcohols in a Jatropa Methyl Ester Fuelled Compression Ignition Engine 1219-1226
15. **Feng Songbao and Sun Linhua**, Statistical and Spatial Analyses of Zinc Concentrations in the Shallow Groundwater of Urban Area and Their Implications on Environmental Background Establishment 1227-1233
16. **Xie Ying chuan and Liu Chang ling**, Construction Dust Emission Features and Management and Control Measures-A Case Study of Zhengzhou City, Henan Province 1235-1241
17. **Preeja P. Thattil and A. Leema Rose**, Photodegradation of Congo Red Dye Via Simple and Effective Air Oxidation Using Copper(II) Chloride and Sunlight 1243-1248
18. **Anwar Ma'ruf, M. Agus Salim Al Fathoni, Agus Mulyadi Purnawanto and Rina Asih Kusumajati**, Development of Hybrid Membrane from Clay/TiO₂-PVA for Batik Wastewater Treatment 1249-1255
19. **Hu Zhaoguang, Ma Xiaorui and Shan Wei**, Health Cost of Dust Pollution of Architectural Engineering Construction in Construction Site: Evidence from China 1257-1263
20. **K. Balasubramanian and K. Purushothaman**, Performance, Emission and Combustion Characteristics of Safflower, Neem and Corn Biodiesels Fuelled in a CI Engine 1265-1273
21. **Bos Ariadi Muis**, Impact of Land Use Change on Hydrological Response of Krueng Aceh Watershed in Aceh Province, Indonesia 1275-1281
22. **Jialei Li, Zhifei Li, Jun Xie, Guangjun Wang, Kai Zhang, Ermeng Yu, Wangbao Gong and Jingjing Tian**, Structures and Anaerobic Metabolic Activity of Attached Microbial Communities During Biofilm Formation in Aquaculture Systems 1283-1291
23. **Yating Huang**, Environmental Pollution Caused by the Transportation Industry and Influencing Factors of Carbon Emission: A Case Study of Jiangxi Province, China 1293-1298

24. **Monangi Murali, R. Srinivasa Rao and Priya Darshini Pradhan**, Some Studies on the Removal of Chromium from Aqueous Solutions by an Adsorbent Obtained from *Terminalia chebula* 1299-1304
25. **Jannathul Firdous, V. Bharathi, S. Durga Devi and J. Jayachitra**, Enzymatic Analysis and Effect of Vermicompost Production from Banana Leaves Waste Using Epigeic Earthworm *Eudrillus euginea* 1305-1311
26. **Ahmed Hasson, Abdulhussien Alaskary and Muhsin Jweeg**, Soil Organic Carbon and Silt-Clay Relationships in the Soil Orders of Northern Agriculture Region (NAR), Western Australia 1313-1318
27. **Wenting Zhang, Yifeng Liu and Yingmei Xu**, A Feasible and Green Approach for Developing Hydro gen Energy 1319-1323
28. **Anushree, Madhu Kumari and Kumar Suranjit Prasad**, Guar Gum Hydrogel Beads for Defluoridation from Aqueous Solution: Kinetic and Thermodynamic Study 1325-1331
29. **G. Srinivas Reddy, K. Vinay Kumar Reddy, B. Linga Reddy and B. Sreenivasa Reddy**, Natural Background Gamma Radiation Levels in the Environs of Proposed Petro-chemical Industry Near Jadcherla, Telangana State, India 1333-1338
30. **Chen Yuling, Wang Chengde, Wu Baoguo and Liu Jiancheng**, Study on Site Quality Assessment of Afforestation Land Based on GA-RBF Neural Network 1339-1347
31. **Chengde Wang, Baoguo Wu, Yuling Chen and Yan Qi**, Development of Crown Profile Models for Chinese Fir Using Non-linear Mixed-Effects Modelling 1349-1361
32. **Priyanka Priyadarshini and Chandan Sahu**, Comparative Assessment of Biochemical Parameters of Plants in Industrial and Non-Industrial Areas of Western Odisha, India 1363-1368
33. **Mingzhao Shen, Meiya Zhu, Jintao Sun, Yunlong Zhang, Rong Li, Fang Niu, Shiyong Li, Xiao Mi and Guoting Li**, Comparison for the Adsorption of Catechol and Hydroquinone on Activated Carbon Fibre: Kinetic Analysis 1369-1374
34. **Chengcai Huang, Rui Qin, Linli Zhang, Muqing Qiu and Linfa Bao**, Adsorption Kinetics and Isotherms of Cr(VI) Ions in Aqueous Solution by Biochar Derived from *Torreya grandis* Nutshell 1375-1380
35. **Miftahul Khair Kadim, Nuralim Pasingi and Sulastri Arsad**, Horizontal Distribution of Chlorophyll-a in the Gorontalo Bay 1381-1385
36. **G. V. Mali**, Toxicological Study of Bifenthrin and its Metabolites on Earthworm (*Eisenia fetida*) 1387-1391
37. **S.G. Antony Godson and S. Gajalakshmi**, High Rate Vermicomposting of Coral Vine by Employing Three Epigeic Earthworm Species 1393-1397
38. **Taufiq Ihsan, Tivany Edwin and Rahma Desri Yanti**, The Effect of Sublethal Exposure of Chlorpyrifos to Nile Tilapia (*Oreochromis niloticus*): Case Study of Twin Lakes of West Sumatra, Indonesia 1399-1403
39. **Emetere M.E., Sanni S.E. and Okoro E.E.**, A Project Design for Air Pollution Studies Over Bobodioulasso-Burkina Faso 1405-1414
40. **Juanqin Xue, Xiande Jing, Shudi Hu, Yuhong Tian, Yonghui Song and Xinzhe Lan**, Process Optimization for the Preparation of Activated Coke from Industrial Waste Using Response Surface Methodology 1415-1421
41. **Khayan Khayan-Taufik Anwar-Slamet Wardoyo**, Plumbum (Pb) in Rainwater in West Kalimantan: Impact of Plumbum (Pb) in Community Blood 1423-1427
42. **Slamet Isworo and Poerna Sri Oetari**, Assessment of Ecological Status in Seawater Around Legon Bajak Port Development Plan, Karimunjawa, Central Java, Indonesia 1429-1446
43. **Kun Xiao and Jingdong Zhang**, Policy Study on Adjustments to Electric Power Prices for China's Air Pollution Abatement 1447-1453
44. **Sucheta Yadav and Subroto Dutta**, Evaluation of Organophosphorus Pesticide Residue in Cotton of Tijara Tehsil, Alwar, Rajasthan 1455-1458
45. **N. Natarajan, V. Vivekk Velusamy, S. Nishar Ahamed, M. Imayaprakash and S. Dineshkumar**, Metal Concentration of Water of Amaravathi and Thirumoorthy Reservoirs in Tamil Nadu, India 1459-1462
46. **Aurora S. Nakpil, Emmanuel S. Bajaand Paul Mark B. Medina**, Mutagenicity of Bulk, Aqueous and Organic Partitions of Air Particulate Matter in Differentially Ventilated Wards in a Public Urban Hospital 1463-1469
47. **Amit Arora, Rohit Mehra, Rajeev Kumar Garg and Anand K. Tyagi**, Assessment of Annual Effective Dose from the Indoor Radon in Bathinda District of Punjab in India 1471-1478

**The Journal
is
Currently
Abstracted
and
Indexed
in:**

International Scientific Indexing (UAE) with Impact Factor 2.236 (2018)

NAAS Rating of the Journal (2019) = 3.85

Scopus®, SJR (0.146) 2018

Index Copernicus (2016) = 109.45

El Compendex of Elsevier

Indian Science Abstracts,
New Delhi, India

Chemical Abstracts, U.S.A.

Elsevier Bibliographic
Databases

Pollution Abstracts, U.S.A.

Zoological Records

Paryavaran Abstract,
New Delhi, India

Indian Citation Index (ICI)

Electronic Social and Science
Citation Index (ESSCI)

EBSCO: Environment Index™

Zetoc

Google Scholar

ProQuest, U.K.

J-Gate

Environment Abstract, U.S.A.

British Library

Centre for Research Libraries

WorldCat (OCLC)

JournalSeek

Connect Journals (India)

CSA: Environmental Sciences and Pollution Management

Research Bible (Japan)

Indian Science

Geobase

Elektronische
Zeitschriftenbibliothek (EZB)

SHERPA/RoMEO

Directory of Science

CNKI Scholar (China National
Knowledge Infrastructure)

Access to Global Online Research in Agriculture (AGORA)

AGRIS (UN-FAO)

Full papers are available on the Journal's Website:
www.neptjournal.com

UDL-EDGE (Malaysia) Products like *i*-Journals, *i*-Focus and *i*-Future

The journal is also included in the UGC CARE list of journals in India

www.neptjournal.com

Nature Environment and Pollution Technology

EDITORS

Dr. P. K. Goel

Former Head, Deptt. of Pollution Studies
Yashwantrao Chavan College of Science
Vidyanagar, Karad-415 124
Maharashtra, India

Dr. K. P. Sharma

Former Professor, Ecology Lab, Deptt. of Botany
University of Rajasthan
Jaipur-302 004, India
Rajasthan, India

Manager Operations: Mrs. Apurva Goel Garg, C-102, Building No. 12, Swarna CGHS, Beverly Park, Kanakia, Mira Road (E)-401107, Distt. Thane, Maharashtra, India (**E-mail: operations@neptjournal.com**)

Business Manager: Mrs. Tara P. Goel, Technoscience Publications, A-504, Bliss Avenue, Balewadi, Pune-411 045, Maharashtra, India (**E-mail: contact@neptjournal.com**)

All correspondence regarding publication of papers in the journal must be made only by e-mail (contact@neptjournal.com) to the Editor

EDITORIAL ADVISORY BOARD

1. **Dr. Prof. Malay Chaudhury**, Department of Civil Engineering, Universiti Teknologi PETRONAS, Malaysia
2. **Dr. Saikat Kumar Basu**, University of Lethbridge, Lethbridge AB, Canada
3. **Dr. Sudip Datta Banik**, Department of Human Ecology, Cinvestav-IPN Merida, Yucatan, Mexico
4. **Dr. Elsayed Elsayed Hafez**, Deptt. of Molecular Plant Pathology, Arid Land Institute, Egypt
5. **Dr. Dilip Nandwani**, College of Agriculture, Human & Natural Sciences, Tennessee State Univ., Nashville, TN, USA
6. **Dr. Ibrahim Umaru**, Department of Economics, Nasarawa State University, Keffi, Nigeria
7. **Dr. Tri Nguyen-Quang**, Department of Engineering Agricultural Campus, Dalhousie University, Canada
8. **Dr. Hoang Anh Tuan**, Deptt. of Science and Technology, Ho Chi Minh City University of Transport, Vietnam
9. **Mr. Shun-Chung Lee**, Deptt. of Resources Engineering, National Cheng Kung University, Tainan City, Taiwan
10. **Samir Kumar Khanal**, Deptt. of Molecular Biosciences & Bioengineering, University of Hawaii, Honolulu, Hawaii
11. **Dr. Sang-Bing Tsai**, Zhongshan Institute, University of Electronic Science and Technology, China
12. **Dr. Zawawi Bin Daud**, Faculty of Civil and Environmental Engg., Universiti Tun Hussein Onn Malaysia, Johor, Malaysia
13. **Dr. Srijan Aggarwal**, Civil and Environmental Engg. University of Alaska, Fairbanks, USA
14. **Dr. M. I. Zuberi**, Department of Environmental Science, Ambo University, Ambo, Ethiopia
15. **Dr. Prof. A.B. Gupta**, Dept. of Civil Engineering, MREC, Jaipur, India
16. **Dr. B. Akbar John**, Kulliyah of Science, International Islamic University, Kuantan, Pahang, Malaysia
17. **Dr. Bing Jie Ni**, Advanced Water Management Centre, The University of Queensland, Australia
18. **Dr. Prof. S. Krishnamoorthy**, National Institute of Technology, Tiruchirappally, India
19. **Dr. Prof. (Mrs.) Madhoolika Agarwal**, Dept. of Botany, B.H.U., Varanasi, India
20. **Dr. Anthony Horton**, Envirocarb Pty Ltd., Australia
21. **Dr. C. Stella**, School of Marine Sciences, Alagappa University, Thondi -623409, Tamil Nadu, India
22. **Dr. Ahmed Jalal Khan Chowdhury**, International Islamic University, Kuantan, Pahang Darul Makmur, Malaysia
23. **Dr. Prof. M.P. Sinha**, Dumka University, Dumka, India
24. **Dr. G.R. Pathade**, H.V. Desai College, Pune, India
25. **Dr. Hossam Adel Zaqoot**, Ministry of Environmental Affairs, Ramallah, Palestine
26. **Dr. Riccardo Buccolieri**, Deptt. of Atmospheric Physics, University of Salento-Dipartimento di Scienze e Tecnologie Biologiche ed Ambientali Complesso Ecotekne-Palazzina M S.P. 6 Lecce-Monteroni, Lecce, Italy
27. **Dr. James J. Newton**, Environmental Program Manager, 701 S. Walnut St. Milford, DE 19963, USA
28. **Prof. Subhashini Sharma**, Dept. of Zoology, University of Rajasthan, Jaipur, India
29. **Dr. Murat Eyvaz**, Department of Environmental Engg. Gebze Inst. of Technology, Gebze-Kocaeli, Turkey
30. **Dr. Zhihui Liu**, School of Resources and Environment Science, Xinjiang University, Urumqi, China
31. **Claudio M. Amescua García**, Department of Publications, Centro de Ciencias de la Atmósfera, Universidad Nacional Autónoma de México
32. **Dr. D. R. Khanna**, Gurukul Kangri Vishwavidyalaya, Haridwar, India
33. **Dr. S. Dawood Sharief**, Dept. of Zoology, The New College, Chennai, T. N., India
34. **Dr. Amit Arora**, Department of Chemical Engineering, Shaheed Bhagat Singh State Technical Campus, Ferozepur -152004, Punjab, India
35. **Dr. Xianyong Meng**, Xinjiang Inst. of Ecology and Geography, Chinese Academy of Sciences, Urumqi, China
36. **Dr. Sandra Gómez-Arroyo**, Centre of Atmospheric Sciences, National Autonomous University, Mexico
37. **Dr. Nirmal Kumar, J. I.**, ISTAR, Vallabh Vidyanagar, Gujarat, India
38. **Dr. Wen Zhang**, Deptt. of Civil and Environmental Engineering, New Jersey Institute of Technology, USA



3D Simulation Model of Tidal, Internal Mixing and Turbulent Kinetic Energy of Palu Bay

Sabhan*†, Alan Frendy Koropitan**, Mulia Purba*** and Widodo Setiyo Pranowo****

*Marine Science, Postgraduate Study Program, IPB University/Department of Physics, Faculty of Mathematical and Natural Sciences, Tadulako University, Indonesia, Perumahan Bumi Mallantan Rezkitia Blok C2/1, Jl Roviega, Tondo, Palu City, Central Sulawesi, Indonesia

**Department of Marine Science and Technology, Faculty of Fisheries and Marine Science, IPB University, Jl.Raya Dramaga, Bogor 16680, West Java, Indonesia

***Department of Marine Science and Technology, Faculty of Fisheries and Marine Science, IPB University, Jl.Raya Dramaga, Bogor 16680, West Java, Indonesia

****Department of Hydrography, Indonesian Naval Postgraduate School (STTAL)/Researcher at Marine Research Center, Indonesian Ministry of Marine Affairs & Fisheries, Indonesia

† Corresponding author: Sabhan

Nat. Env. & Poll. Tech.
Website: www.neptjournal.com

Received: 18-02-2019

Accepted: 22-05-2019

Key Words:

3D simulation model
Tidal simulation models
Turbulent kinetic energy
Palu bay

ABSTRACT

The tidal simulation models and internal mixing were constructed using finite volume method to simulate diurnal tide (K1) constituent and semidiurnal tide (M2) constituent, the mixing level model using General Ocean Turbulence Model (GOTM). Tidal elevation amplitude of the model K1 ranges from 19.27 to 19.31 cm, high tidal amplitude at the point near the mouth of the Palu River, low tidal amplitude at the open boundary of the model that leads to the Makassar Strait. The amplitude range by M2 tidal constituents is 55.55-55.75 cm, high tidal amplitude at the mouth of Palu Bay and the end of Palu Bay and the slope area. The tidal current of the K1 constituent strengthens at the open boundary of the model and weakens into Palu Bay, which then undergoes strengthening near the mouth of the Palu River which experiences extreme siltation, tidal currents strengthen in the mouth area of Palu Bay and the tip of Palu Bay is caused by the tidal constituent propagation M2 and also the slope of the area along the coast of Palu Bay. Bottom Ekman layer which is caused by K1 constituent can reach 11 meters while the M2 constituent reaches 7 meters. In the slope area, are also found the buoyancy frequency with the order of 10-5 which is at a depth of 75-150 m. Kinetic energy in Palu Bay stands at $O(10^{-5}-10^{-3})$ with high kinetic energy around rough topography in the order of 10^{-3} around the mouth of Palu Bay.

INTRODUCTION

Palu Bay is administratively located in Donggala Regency, Palu City of Central Sulawesi Province. The position of Palu Bay is facing north with the bay mouth leading to the Makassar Strait, which is the main line of Indonesian Throughflow. The circulation of Palu Bay as semi-closed waters is strongly influenced by tides. Palu Bay has a unique morphology with a high depth degradation at a distance of 2000 meters from the coast, which can reach the depth of 300 meters or a slope of 25%. It is very potential for the occurrence of internal mixing on the slope. The process of seawater circulation is important to detect areas that have robust internal mixing in the management of the marine environment.

The depth of 450 meters in the mouth of Palu Bay allows the potential of Arlindo flow to affect the circulation in Palu Bay. At the end of the Palu Bay there is a major river that flows with an average discharge of $83.6 \text{ m}^3/\text{s}$ with a

maximum discharge of $423 \text{ m}^3/\text{s}$ which is possible to carry sediment of 0.42244 kg/s/m (Rusdin et al. 2014) from the upstream which are intensely experiencing land clearing for new agriculture land. This high level of discharge enables the potential nutrient transport into the Palu Bay.

Research on the 2D tidal circulation model in the bay area has been conducted by Carbajal & Backhaus (1998) with the semi-implicit method showing that tidal energy dissipation is found very strong in the delta. A 3D model study conducted by Koropitan & Ikeda (2008) with a larger scale on the Java Sea shows that the mixing is dominant by the K1 tidal constituent. Pichon & Corréard (2006) modelled internal tides using a spring tidal constituent of daily tidal, showing that the vertical distribution of internal tidal amplitude is related to baroclinic propagations. Kartadikaria et al. (2011) conducted research in the Indonesian Sea using POM with $(1/36)^\circ$ resolution showing that vertical

mixing increased due to the effects of baroclinic pairs and increased horizontal mixing with the presence of barotropic tidal movements. Increased mixing has a role in eliminating high salinity water masses from the Pacific Ocean. Robertson & Field (2005) stated that in the Indonesian Sea significant M2 values of baroclinic tides were found on slope and rough topography, especially in the strait.

Tidal circulation modelling is an essential step to apply as it is being the main cause of internal mixing in the slope that occurs by reflection of M2 internal waves (Nash et al. 2004). Strong internal mixing indicates high primary productivity, although the internal mixing has insignificant contribution (10% to 25%) to primary productivity (Schafstall et al. 2010). Internal mixing causes nutrients to be mixed in the sediment so that they enter the upper water column which is then utilized by phytoplankton which is characterized by increased chlorophyll-*a* (McGowan & Hayward 1978, Pingree & Mardell 1981). Internal mixing is a mechanism for lifting nutrients to the surface in increasing primary productivity (Sharples et al. 2001, Holt et al. 2012). In the previous study, the focus was on regional areas, while this study focused on more detailed areas with a grid size of 100 m so that small degradation could be clearly identified.

This research simulates hydrodynamics in 3D, which

is driven by wind and tides by using numerical models of finite volume. The result is verified by using measurements of currents and tides. This study investigates significant tidal constituents and its role to the mixing level. The total turbulent kinetic energy and dissipation in the domain model area are also computed.

MATERIALS AND METHODS

Description of the study sites: This research was conducted in Palu Bay, Central Sulawesi Province, Indonesia. Palu Bay is located on the Arlindo line in the Makassar Strait and the input of the Palu River is large during the rainy season. Rainfall patterns in the Palu bay fall into the category of Local Rain Patterns with only one (unimodal) peak season with an average annual rainfall of 547 mm, the lowest in all of Indonesia with the peak of the rainy season in June (Bayong 1999). Palu Bay provides environmental quality with human activities as the central government of Central Sulawesi Province (Fig. 1).

Hydrodynamic model, turbulent and validation: Hydrodynamic modelling in Palu Bay using 3D MOHID uses a finite volume approach to discretize equations (Neves 2007). MOHID 3D is a hydrodynamic model based on the Navier-Stokes equation with a Boussinesq and hydrostat-




Fig. 1: the location of Teluk Palu, Central Sulawesi Province, Indonesia.

ic approach. 3D MOHID is formulated in a volume up to vertical discretization approach using sigma coordinates (Neves 2007). The open boundary of the model which is the generator input point for the tidal mouth model of Palu Bay leading to the Makassar Strait uses tidal constituents from FES2004. The study was carried out in Palu Bay (Fig. 1) by building a model that covers the entire water body of the Palu Bay. Turbulence modelling was carried out using a module (GOTM) which has been integrated with MOHID (Baumert et al. 2005) using the κ - ϵ model approach (Versteeg & Malalasekera 2007).

Horizontal turbulence is calculated using a Smagorinsky turbulence model based on grid size and local derivative:

$$v_H = \Delta x \Delta y \sqrt{\left(\frac{\partial u}{\partial x}\right)^2 + \left(\frac{\partial v}{\partial y}\right)^2 + \frac{1}{2} \left(\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y}\right)} \quad \dots(1)$$

Equation (1) in MOHID has an expression in the form (Marín et al. 2013):

$$v_H = \text{Horcon} \times \Delta x \times \Delta y \sqrt{\left(\frac{\partial u}{\partial x}\right)^2 + \left(\frac{\partial v}{\partial y}\right)^2 + \frac{1}{2} \left(\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y}\right)} \quad \dots(2)$$

Where, $\text{Horcon} = 2.5$ (Marín et al. 2013), is a dimensionless calibration constant.

Turbulence parameters in the form of vertical eddy diffusion by internal waves are calculated by the following equation (Gargett & Holloway 1984):

$$K_z = \frac{R_f}{1 - R_f} \frac{\epsilon}{N^2} \quad \dots(3)$$

Where, R_f is the Richardson number flux (0.20). Vertical displacement is obtained from the equation (Wan et al. 2018):

$$APE = \frac{g \rho'}{2 \rho_c N^2} \quad \dots(4)$$

Where, g is gravity acceleration, ρ is the perturbation density, ρ_c is the reference density, and N is the buoyancy frequency.

The grid cell size is $0.001^\circ \times 0.001^\circ$. As many as 14 main constituents of tidal (Mf, Mm, Msqm, Mtm, O1, P1, Q1, K1, M2, K2, 2N2, N2, S2 and M4) are used as tidal generators at the open boundary, and these constituents are derived from the tide model FES2004 global tide (Lyard et al. 2006). Tidal data were also obtained from the Pantoloan Port station operated by the Geospatial Information Agency (BIG). Bathymetry data were obtained from the Navy Base (Lanal) Palu with a grid of 20×20 meters (not accommodating bathymetry changes by the 2018 earthquake and tsunami incident).

Model Analysis: Characteristics of water level by tides and tidal currents resulting from the model based on the tidal harmonic constants of the two dominant tidal constituents (M2 and K1) were analysed with T_TIDE program (Pawlowicz et al. 2002).

RESULTS

Validation of model: The model was run for 30 days and validated with the results of tidal measurements from the Geospatial Information Agency (BIG) in Pantoloan Port for January (Table 1), the velocity of the model results was validated with the results of measurements at Donggala Beach (Fig. 2). K1 tidal constituent has the greatest contribution to the occurrence of discrepancies between the results of observation (Fig. 2) and results of model with the difference in amplitude and phase respectively 2.41 cm and -12.17 (Table 1).

Tidal Simulation

Tidal constituent K1: The amplitude, phase, and tidal currents of the K1 constituent are calculated for each 5 grids. Co-amplitude, co-phase and elliptical tidal currents of the K1 constituent are shown in Fig. 3a. The co-amplitude pattern and co-phase show a high tidal amplitude pattern at the end of the bay compared to the mouth of Palu Bay, while the phase propagates from the bay mouth who experience resonance in the middle of the bay due to the reflection of the wave returning from the end of the bay. The tidal eleva-

Table 1: Comparison of amplitude and phase between data from BIG and model results.

Constituent	Amplitudes (cm)			Phase (°)		
	Model	BIG Data	ΔH	Model	BIG Data	$\Delta HO1$
$\Delta HO1$	15.48	16.35	-0.87	128.09	131.57	-3.48
K1	19.06	21.47	-2.41	138	150.17	-12.17
M2	54.38	55.1	-0.72	284.09	282.38	1.71
S2	42.97	42.05	0.92	337.83	332.74	5.09




Fig. 2: Zonal and meridional current velocity plots between tidal currents of observation results (red) and model results (blue).




Fig. 3: Co-amplitude (cm) and Co-phase ($^{\circ}$) (a) tidal constituent of K1, (b) tidal constituent of M2, amplitude (solid line) co-phase (dashed line).

tion amplitude of the K1 model results ranged from 19.27 to 19.31 cm high amplitude at points close to the mouth of the Palu River and low at the open boundary of the model leading to the Makassar Strait.

Co-phase model results show that the tidal wave propagates from the open boundary of the model in the Makassar Strait entering the bay until it reaches the end of the bay at the Palu river mouth with tidal co-phase from 150.3–150.42°.

Tidal current constituent K1 is presented from the model results for the surface (Fig. 4b) and bottom (Fig. 4a) of the waters in Palu Bay. The pattern of velocity of tidal currents of the K1 constituent strengthens at the open boundary of the model and weakens into the Palu Bay, which then undergoes strengthening at a certain point near the estuary which experiences extreme silting, the pattern between the speed on the surface and the bottom of the waters resembles only the 25% stronger surface compared to velocity at the bottom.

The average current velocity of the K1 constituent model results shows that there was a fairly strong current heading north near the east coast of the bay with a velocity of about 0.5 m/s in January 2016, the same as the research conducted by Sabhan et al. (2019). A strong K1 constituent

velocity is also found in the area near the mouth of the Palu Bay in a shallow topographic area (Fig. 6a).

Tidal constituent M2: Elliptical parameters of tidal currents by tidal constituents M2 was calculated for every 10 grids. Co-tidal is shown in Fig. 3b, referring that the M2 constituent fluctuates from the open boundary of the model at the mouth of Palu Bay then strengthens at the mouth of the Palu River at the end of the bay so that the co-amplitude is low in the centre of the bay, the amplitude range of M2 tidal constituents are 55.55 – 55.75 cm. Fig. 3b shows that the high amplitude in the bay mouth area, at the bay end and the slope areas which are close to the beach.

The velocity of the tidal current represented by the tidal current ellipse by the M2 constituent shows the same pattern between the surface and the bottom of the waters, but at the latter is weakened due to friction with the bottom of the waters (Fig. 5a). Tidal currents strengthened in the bay mouth area and the gulf tip (Fig. 5b) by the M2 tidal constituent propagation and the slope areas along the coast of Palu Bay. That corresponds to the results of research on the Hawaiian Sea backs that experienced siltation (Merrifield & Holloway 2002). Because Palu Bay only opens towards the Makassar Strait, the tidal propagation of M2 constituents comes from the Makassar Strait.




Fig. 4: Tidal current ellipses for K1 constituent (a) bottom (b) Surface.




Fig. 5: Tidal current ellipses for M2 constituent (a) bottom, (b) Surface.

A strong M2 constituent velocity was also found in the area near the estuary of Palu Bay in a shallow topographical area (Fig. 6b). The M2 constituent pattern is almost the same as the K1 constituent pattern but the M2 constituent is stronger than the K1 constituent. Besides that, it is found that a fairly strong average current velocity reaching 1 m/s near the Palu River estuary is occurring in an area with low depth. Robertson & Field (2005) found that in the Indonesian Sea strong M2 baroclinic tides were present on the slope and rough topography especially in the strait.

Turbulent Models

Buoyancy frequency: The results of the model show that the buoyancy frequency (Brunt-Väisälä frequency) is in order (10^{-1} - 10^{-9}) s^{-2} . The cross section intersecting the y axis that passes through the end of Palu Bay (estuary of Palu River) to the mouth of Palu Bay (Makassar Strait) shows that the frequency of buoyancy is high in the bay area, which is close to the river mouth, where the input of high river water forms a strong stratification by difference density. This result corresponds to the results of the study by Hordoir et al. (2018) which states that high buoyancy frequency values are present under strong stratification conditions. In the slope area also found areas with the order 10^{-5} which are at a depth of 75-150 m with a depth of

about 25 m from the bottom of the waters (Fig. 11a), this occurs because the slope area is a region with high friction (Wunsch 1968).

Turbulent dissipation: Fig. 10a shows the turbulent dissipation distribution pattern near the bottom in the Palu Bay. Fig. 11b shows that turbulent dissipation in the slope area is at $O(10^{-5}$ - $10^{-4})$, whereas in areas with rough topography there is a turbulent dissipation area that is small, but higher one to two orders from the surrounding environment. These results are consistent with the results of the model conducted by Nikurashin et al. (2013) which states that in the absence of rough topography, internal turbulent mixing will decrease. As for the ocean with small tidal currents, internal wave friction is the dominant cause of turbulent dissipation (Ledwell et al. 2000, Jayne & St Laurent 2001).

Eddy diffusivity: Vertical slice of vertical eddy diffusivity on the $y = 33$ axis (Fig. 11c) has a fairly wide order with (10^0 - 10^{-9}), areas that have high eddy diffusivity in areas close to coarse topography with $O(10^{-3}$ - $10^{-2})$ and the surface area, in the mouth area of the Bay found a high diffusivity area with first order where the incoming tidal currents hit the rough topography. According to the results of the study (Toole et al. 1994), increased dissipation was




Fig. 6: The average velocity vector of (a) K1 constituent, (b) M2 constituent.

observed in the areas near steep boundaries (eddy diffusivity $>10^{-4}$ square meters per second).

Bottom friction velocity: Bottom friction velocity was calculated using equations (Zhurbas et al. 2018):

$$u^* = (\tau/\rho)^{0.5} \quad \dots(5)$$

$$\tau = \rho C_d u^2 \quad \dots(6)$$

Where, τ is the basic shear stress, ρ is the density of seawater. Fig. 8, the bottom friction velocity distribution for the tidal K1 and M2 constituents shows that the basic friction velocity of K1 constituent (Fig. 8a) has increased at a point where there is a rough topography, but is rapidly weakened, while the M2 constituent (Fig. 8b) speeds up from the bay mouth until reach the middle of the bay. High velocity is also found at the end of the bay due to silting or slope plus the flow of the river at the mouth of the Palu River. Sea threshold topography causes constriction, increasing the velocity of basic friction, this implies an increase in mixing on the other side of the sea threshold topography (Xing & Davies 2009).

Delta Ekman layer: Delta Ekman layer is calculated using equation (He & Weisberg 2002):

$$\delta = \frac{cu^*}{\sigma + f} \quad \dots(7)$$

Where, $c = 0.2-0.4$, σ is the tidal frequency and f is the tidal parameter. The M2 tidal frequency for Palu Bay is 2 times the frequency of the coriolis parameter so that the Ekman layer for the tidal constituent of M2 becomes: $\delta = \frac{cu^*}{3f}$.

Fig. 9 shows that the Ekman layer by the K1 constituent (Fig. 9a) can reach 11 meters, while the M2 constituent (Fig. 9b) reaches 7 meters. The point undergoes a high layer Ekman on the rough topography of the north side of the mouth of Palu Bay. But what is effective in mixing is in the slope section where the sea is getting shallow and the layer Ekman is getting bigger so that the mixed layer from the base is getting higher, in the next study this area has the potential to supply nutrients to the water column.

Energy flux is calculated using the equation:

$$(E_x, E_y) = \frac{\rho_0}{T} \int_0^T \int_{-h}^{\eta} (u, v) \left[g\eta + \frac{u^2 + v^2}{2} \right] dz dt \quad \dots(8)$$

Where, E_x and E_y are each flux energy of the constituent vector x and y , (u, v) is the vector of horizontal velocity,




Fig. 7: Energy flux of Palu Bay by (a) tidal constituent K1, (b) tidal constituent M2.




Fig. 8: Bottom friction velocity by (a) tidal constituent of K1, (b) tidal constituent of M2.




Fig. 9: Bottom Ekman layer by (a) Tidal constituent of K1, and (b) Tidal constituent of M2.




Fig. 10 (a): Turbulent dissipation (ϵ) (b) turbulent kinetic energy Palu Bay.




Fig. 11: Vertical slice of (a) buoyancy frequency, (b) eddy diffusivity, (c) turbulent dissipation, (d) turbulent kinetic energy, (e) vertical displacement in the waters of Palu Bay.

“ η ” is the elevation of the water level, “ g ” is acceleration of gravity, “ h ” is the depth of seawater, “ T ” is the period of tides, ρ_0 density of seawater. The results of the model show that M2 kinetic energy of turbulent (Fig. 7b) is greater than K1 (Fig. 7a) with the dominant energy vector to the north with large on the right side of Palu Bay.

Turbulent kinetic energy: The kinetic energy in Palu Bay is at O (10^{-5} - 10^{-3}) with high kinetic energy found around the rough topography in the 10^{-3} order around the mouth of Palu Bay (Fig. 11d), the roughness effect tends to lower the average velocity distribution and increase turbulent kinetic energy within the boundary layer (Rachman et al. 2011).

CONCLUSION

The tidal elevation amplitude of the model K1 ranges from 19.27 to 19.31 cm, high tidal amplitude at the point near the mouth of the Palu River, and low tidal amplitude at the open boundary of the model that leads to the Makassar Strait. The amplitude range of M2 tidal constituents is 55.55 - 55.75 cm, high tidal amplitude at the mouth of Palu Bay and the end of Palu Bay and the slope area. The tidal current of the K1 constituent is strengthened at the open boundary of the model and weakened when it enters Palu Bay which then amplify near the mouth of the Palu River because of its extreme siltation. Tidal currents strengthen in the mouth area and the tip of Palu Bay by propagation of the tidal constituent M2 and the area in the slope along the coast of Palu Bay. The basic Ekman layer by the K1 constituent can reach 11 meters while the M2 constituent reaches 7 meters. In the slope area also found the buoyancy frequency with the order of 10^{-5} which is at a depth of 75-150 m. The kinetic energy in Palu Bay stands at O (10^{-5} - 10^{-3}) with the highest kinetic energy found in the rough topography of the mouth of Palu Bay in the order of 10^{-3} .

REFERENCES

- Baumert, H.Z., Simpson, J. and Sündermann, J. 2005. *Marine Turbulence: Theories, Observations, and Models*. Cambridge University Press.
- Bayong T. 1999. *Klimatologi Umum*. Penerbit ITB Bandung.
- Carbajal, N. and Backhaus, J. 1998. Simulation of tides, residual flow and energy budget in the Gulf of California. *Oceanologica Acta.*, 21(3): 429-446.
- Fortin, W.F., Holbrook, W.S. and Schmitt RW. 2016. Mapping turbulent diffusivity associated with oceanic internal lee waves offshore Costa Rica. *Ocean Sci.*, 12: 601-612.
- Gargett, A.E. and Holloway, G. 1984. Dissipation and diffusion by internal wave breaking. *Journal of Marine Research*, 42(1):15-27.
- He, R. and Weisberg, R.H. 2002. Tides on the west Florida shelf. *J. of Physical Oceanography*, 32(12): 3455-3473.
- Holt, J., Butenschon M., Wakelin S., Artioli Y. and Allen J. 2012. Oceanic controls on the primary production of the northwest European continental shelf: Model experiments under recent past conditions and a potential future scenario. *Biogeosciences*, 9: 97-117.
- Hordoir, R., Höglund, A., Pemberton, P. and Schimanke, S. 2018. Sensitivity of the overturning circulation of the Baltic Sea to climate change, a numerical experiment. *Climate Dynamics*, 50(3-4): 1425-1437.
- Jayne, S.R. and St Laurent, L.C. 2001. Parameterizing tidal dissipation over rough topography. *Geophysical Research Letters*, 28(5): 811-814.
- Kartadikaria, A., Miyazawa, Y., Varlamov, S. and Nadaoka, K. 2011. Ocean circulation for the Indonesian seas driven by tides and atmospheric forcings: Comparison to observational data. *Journal of Geophysical Research: Oceans*, 116(C9).
- Koropitan, A.F. and Ikeda, M. 2008. Three-dimensional modelling of tidal circulation and mixing over the Java Sea. *Journal of Oceanography*, 64(1): 61-80.
- Ledwell, J., Montgomery, E., Polzin, K., Laurent, L.S., Schmitt, R. and Toole, J. 2000 Evidence for enhanced mixing over rough topography in the abyssal ocean. *Nature*, 403(6766): 179.
- Lyard, F., Lefevre, F., Letellier, T. and Francis, O. 2006. Modelling the global ocean tides: Modern insights from FES2004. *Ocean Dynamics*, 56(5-6): 394-415.
- Marín, V.H., Tironi, A., Paredes, M.A. and Contreras, M. 2013. Modelling suspended solids in a Northern Chilean Patagonia glacier-fed fjord: GLOF scenarios under climate change conditions. *Ecological Modelling*, 264: 7-16.
- McGowan, J.A. and Hayward, T.L. 1978. Mixing and oceanic productivity. *Deep Sea Research*, 25(9): 771-793.
- Merrifield, M.A. and Holloway, P.E. 2002. Model estimates of M2 internal tide energetics at the Hawaiian Ridge. *Journal of Geophysical Research: Oceans*, 107(C8): 5-1-5-12.
- Nagai, T. and Hibiya, T. 2015. Internal tides and associated vertical mixing in the Indonesian Archipelago. *Journal of Geophysical Research: Oceans*, 120(5): 3373-3390.
- Nash, J.D., Kunze, E., Toole, J.M. and Schmitt, R.W. 2004. Internal tide reflection and turbulent mixing on the continental slope. *Journal of Physical Oceanography*, 34(5): 1117-1134.
- Neves, R. 2007. Numerical models as decision support tools in coastal areas. In: Di Dalam (ed.) *Assessment of the Fate and Effects of Toxic Agents on Water Resources*, Springer, pp. 171-195.
- Nikurashin, M., Vallis, G.K. and Adcroft, A. 2013. Routes to energy dissipation for geostrophic flows in the Southern Ocean. *Nature Geoscience*, 6(1): 48.
- Pawlowicz, R., Beardsley, B. and Lentz, S. 2002. Classical tidal harmonic analysis including error estimates in MATLAB using T_TIDE. *Computers & Geosciences*, 28(8): 929-937.
- Pichon, A. and Corréard, S. 2006. Internal tides modelling in the Bay of Biscay. Comparisons with observations. *Scientia Marina*, 70(S1): 65-88.
- Pingree, R.D. and Mardell, G. 1981. Slope turbulence, internal waves and phytoplankton growth at the Celtic Sea shelf-break. *Phil. Trans. R. Soc. Lond. A.*, 302(1472): 663-682.
- Rachman, T., Suntoyo, S., Sambodho, K., Armono, H.D. and Yusroni, E. 2011. Numerical modelling of turbulent bottom boundary layer over rough bed under irregular waves. *IPTEK The Journal for Technology and Science*, 22(4): 177-189.
- Robertson, R. and Ffield, A. 2005. M2 Baroclinic tides in the Indonesian seas. *Oceanography-Washington Dc-Oceanography Society*, 18(4): 62.
- Rusdin, A., Abu, A., R. Kalawawo, P. 2014. Effect of debit on Palu River based sediment movements. *National Civil Engineering Conference*.
- Sabhan, Koropitan A. F., Purba M., Pranowo W. S., Rusydi M. 2019 Numerical model of ocean currents, sediment transport, and geomorphology due to reclamation planning in Palu Bay. *AES Bioflux* 11(2): 87-96.
- Schafstall, J., Dengler, M., Brandt, P. and Bange, H. 2010. Tidal induced mixing and diapycnal nutrient fluxes in the Mauritanian upwelling region. *Journal of Geophysical Research: Oceans*, 115(C10).

- Sharples, J., Moore, C.M. and Abraham, E.R. 2001. Internal tide dissipation, mixing, and vertical nitrate flux at the shelf edge of NE New Zealand. *Journal of Geophysical Research: Oceans*, 106(C7): 14069-14081.
- Toole, J.M., Schmitt, R.W. and Polzin, K.L. 1994. Estimates of diapycnal mixing in the abyssal ocean. *Science*, 264(5162): 1120-1123.
- Versteeg, H.K. and Malalasekera, W. 2007. *An Introduction to Computational Fluid Dynamics: The Finite Volume Method*. Pearson Education, Pearson Education Limited
- Wunsch, C. 1968. On the propagation of internal waves up a slope. *Deep Sea Research and Oceanographic*, 15: 251-258.
- Xing, J. and Davies, A.M. 2009. Influence of bottom frictional effects in sill regions upon lee wave generation and implications for internal mixing. *Ocean Dynamics*, 59(6): 837-861.
- Zhurbas, V., Väli, G., Golenko, M. and Paka, V. 2018. Variability of bottom friction velocity along the inflow water pathway in the Baltic Sea. *Journal of Marine Systems*, 184: 50-58.